

## 10 mg Warfarin Nomogram

Patients < 70 years old and not recently ill

Day 1 = 1<sup>st</sup> day of warfarin

INR should be done in morning and drug taken in early evening

Baseline INR must be < 1.4, platelets > 50

INR should be done daily starting at Day 3 and continuing until INR ≥ 2.0

Day 3 prescribe dose for Day 3 and 4, but still do INR on Day 4

All patients receive 10 mg on Day 1 and Day 2

Day 3		Day 4	Day 5		Day 6	Day 7
INR	Dose	Dose	INR	Dose	Dose	Dose
< 1.3	15	15	< 2.0	15	15	15
1.3 – 1.4	10	10	2.0 – 3.0	7.5	5	7.5
			3.1 – 3.5	0	5	5
			> 3.5	0	0	2.5
1.5 – 1.6	10	5	< 2.0	7.5	7.5	7.5
1.7 – 1.9	5	5	2.0 – 3.0	5	5	5
			3.1 – 3.5	2.5	2.5	2.5
			> 3.5	0	2.5	2.5
2.0 – 2.2	2.5	2.5	< 2.0	5	5	5
2.3 – 3.0	0	2.5	2.0 – 3.0	2.5	5	2.5
			3.1 – 3.5	0	2.5	0
			> 3.5	0	0	2.5
> 3.0	0	0	< 2.0	2.5	2.5	2.5
			2.0 – 3.0	2.5	0	2.5
			3.1 – 3.5	0	2.5	0
			> 3.5	0	0	2.5

Dr. Phil Wells,  
 Canada Research Chair in Thromboembolic Disease Oct 2001  
 Chief, Division of Haematology, The Ottawa Hospital  
 October 2001



Drugs that can <b>diminish</b> the anticoagulant effect of warfarin	Drugs that can <b>Potentiate</b> the anticoagulant effect of warfarin.
Carbamazepine (Tegretol) Cholestyramine Colestipol Cyclosporine Dicloxacillin Ethanol Griseofulvin Nafcillin Pentobarbital Phenobarbital Phenytoin Primidone Rifabutin Rifampin Sulcralfate	Aspirin Acetaminophen Amiodarone Cefotetan Celecoxib Choral hydrate Cimetidine Ciprofloxacin Clarithromycin Clofibrate Cotrimoxazole Disopyramide Disulfiram Erythromycin Fluconazole Gemfibrozil Isoniazid Itraconazole Ketoconazole Levofloxacin Metolazone Metronidazole Norfloxacin Ofloxacin Omeprazole Propafenone Thyroid hormone trazadone